

Quantum Field Theory I: PHYS 721

Problem Set 1

Overview

This question illustrates why we need quantum field theory, and not just relativistic quantum mechanics.

Question 1

[20]

One way of motivating the free-particle Schrödinger equation is to take the non-relativistic dispersion relation $E = \vec{p}^2/(2m)$ and to make the replacements,

$$E \rightarrow i\hbar \frac{\partial}{\partial t}, \vec{p} \rightarrow -i\hbar \vec{\nabla} \quad (1)$$

so that when the wavefunction ψ satisfies

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \vec{\nabla}^2 \psi, \quad (2)$$

it necessarily has the right energy-momentum relationship.

If we follow the same procedure for the relativistic dispersion relation, $E^2 = |\vec{p}|^2 + m^2$ in natural units, we get the free Klein-Gordon equation

$$-\frac{\partial^2 \psi}{\partial t^2} = -(\vec{\nabla}^2 - m^2)\psi \quad (3)$$

and we are tempted to identify ψ with the wavefunction of a single particle (as we did in the Schrödinger case). Is this so bad?

Unfortunately this doesn't work as a consistent theory—in this question we will start to understand why.

The first mathematical issue is that the Klein-Gordon equation is second order in time derivatives, while the Schrödinger equation is first order. This means we'll need to specify both the state of the quantum system and its rate of change at an initial time to get the time-evolution. But suppose we brush this under the carpet. The second issue is that the Klein-Gordon equation admits solutions of negative energy, $E_p = \pm\sqrt{|\vec{p}|^2 + m^2}$. How do we interpret them?




Figure 1: Rectangular barrier used in Question 1(c).

(a) Show that the current, $j^\mu = N(\psi^* \partial^\mu \psi - \psi \partial^\mu \psi^*)$, is conserved, so that $\partial_\mu j^\mu = 0$, if ψ and ψ^* satisfy the Klein-Gordon equation, where N is an arbitrary constant normalisation. Find N by insisting that the non-relativistic limit agree with the probability current of the Schrödinger equation.

(b) We would like to identify j^0 with the probability density. Show that for free-particle solutions of the Klein-Gordon equation, $\psi = ae^{-ik^\mu x_\mu}$ and $j^0 = (E_k/m)|a|^2$. Notice that the negative energy solutions have negative probability density! In general solutions of the Klein-Gordon equation are not guaranteed to have positive probability density.

Suppose we brush this issue of negative probabilities under the carpet, too. Let's see what happens if we introduce "interactions" in the way we would in the Schrödinger equation, by introducing a potential,

$$i \frac{\partial}{\partial t} \rightarrow i \frac{\partial}{\partial t} - V. \quad (4)$$

Consider the rectangular barrier shown below, in Figure 1. For the rest of this question, assume that we're working in one spatial dimension.

(c) For positive energy waves of energy E incident from the left we have

$$p = \sqrt{E^2 - m^2}, \quad p' = \sqrt{(E - V - m)(E - V + m)}. \quad (5)$$

Using continuity of ψ and $\partial\psi/\partial x$, show that the transmitted fraction is

$$|T|^2 = \left| \cos(ap') - \frac{i}{2} \left(\frac{p}{p'} + \frac{p'}{p} \right) \sin(ap') \right|^{-2}. \quad (6)$$

For (kinetic) energies above the barrier, $E - m > V$, p' is real and $|T| < 1$, which corresponds to some reflection of flux. This is in accord with non-relativistic intuition.

For energies below a relatively low barrier, $E - m < V < E + m$, p' is imaginary and we have tunnelling wavefunctions between $x = 0$ and $x = a$. This is also in accord with non-relativistic intuition.

For a very high barrier, we'd expect to continue to have tunnelling, but with an exponentially decreasing amplitude. In fact we get something completely different: For $V > E + m$ (which implies $V > E - m$ necessarily), p' is real again, and we have propagating solutions between $x = 0$ and $x = a$. The probability density in this region is also negative. This is known as Klein's paradox and it cannot be resolved without realising that the KG equation is a field equation associated with an indefinite number of particles, and not a single-particle wave equation.