

Quantum Field Theory I: PHYS 721

Problem Set 8

Chris Monahan

Overview

The questions in this problem set familiarise you with deriving Feynman rules from an example of an interacting theory, and writing down Feynman diagrams corresponding to time-ordered correlation functions. In many applications, this is a typical way to carry out calculations in QFT—given a Lagrangian, you derive the Feynman rules, write the Feynman diagrams that correspond to that process at a given order in perturbation theory (and then calculate the diagrams). There are two questions.

Question 1

[12]

In our mini-lectures we derived the Feynman rules for $\lambda\phi^4$ theory, which ended up being pretty simple. In fact, it is usually possible to just read off the Feynman rules directly from the Lagrangian. Consider the following Lagrangian¹,

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m^2\phi^2 - \frac{\lambda}{3!}\phi^3,$$

which defines the so-called $\lambda\phi^3$ theory (very imaginatively named, as you can see).

(a) What is the mass dimension of the coupling λ in four spacetime dimensions? Is this coupling relevant, marginal, or irrelevant in four spacetime dimensions? In how many spacetime dimensions is λ dimensionless?

(b) Write out what you expect the relevant position-space Feynman rules to be. You do not need to provide a detailed derivation via Wick's theorem.

(c) Write out what you expect the relevant momentum-space Feynman rules to be. You do not need to provide a detailed derivation.

(d) Draw all the connected diagrams contributing, up to and including $\mathcal{O}(\lambda^2)$, to the:

(i) The two-point function

$$\langle\Omega|T\{\phi(x)\phi(y)\}|\Omega\rangle.$$

(ii) The three-point function

$$\langle\Omega|T\{\phi(x)\phi(y)\phi(z)\}|\Omega\rangle.$$

(iii) The four-point function

$$\langle\Omega|T\{\phi(w)\phi(x)\phi(y)\phi(z)\}|\Omega\rangle.$$

¹Be aware! This is not exactly the same Lagrangian as we considered in class, or Schwartz Section 7.4!

Question 2**[8]**

Consider scalar Yukawa theory, defined by the Lagrangian

$$\mathcal{L} = \partial_\mu \psi^* \partial^\mu \psi - M^2 \psi^* \psi + \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 - g \psi^* \psi \phi,$$

where ψ is a complex scalar field and ϕ is a real scalar field. Scalar Yukawa theory is a toy model for the low energy interactions of hadrons, such as pions and neutrons and protons. In reality, many hadrons are fermions, so the full Yukawa model (in which a scalar field couples to spinor fields) is a more realistic representation of hadron interactions at low energy. As the energy of the process increases, both of these models start to fail and the process must be calculated in the full theory of the strong interaction, quantum chromodynamics (which governs the behaviour of quarks and gluons within hadrons, and the interactions between hadrons).

(a) Draw the leading nontrivial (i.e. do not count the case of no scattering) Feynman diagrams for the following processes, and express the invariant matrix element for each process in terms of Mandelstam variables:

1. $\psi\psi$ scattering, $\psi\psi \rightarrow \psi\psi$;
2. $\psi\psi^*$ scattering, $\psi\psi^* \rightarrow \psi\psi^*$;
3. ϕ pair production, $\psi\psi^* \rightarrow \phi\phi$;
4. $\psi\phi$ scattering, $\psi\phi \rightarrow \psi\phi$.

(b) Draw the leading nontrivial (i.e. do not count the case of no scattering) Feynman diagrams for the following processes:

1. ϕ decay, $\phi \rightarrow \psi\psi^*$ (assuming $m > 2M$);
2. $\phi\phi$ scattering, $\phi\phi \rightarrow \phi\phi$.